

Aprice Hamoi, Vietnam

information for a Smarter Life and Environmental Resilience"



Multi-Layer Optimisation Technique (M-LOT) for **Shallow Water Hydrographic Survey Mapping Using** Satellite-derived Bathymetry Application

First Admiral Dr Najhan MD SAID **National Hydrographic Centre** Port Klang, Malaysia

Sr Dr Rozaimi CHE HASAN Universiti Teknologi Malaysia Kuala Lumpur, Malaysia










22-26 April, Hanoi, Vietnam






Background


Acquisition technique of bathymetric data has evolve from shipborne platform to airborne and even by means of space-borne acquisition







Images retrieved from the NOAA Office of Coast Survey website: https://www.nauticalcharts.noaa.gov.











22–26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"



Background

- SDB is an acquisition method related to a systematic analysis modelling of the recorded light penetration intensity through the water column at different spectral bands (multi/hyperspectral)
- Various factors affected
 - height and direction of the sun
 - clouds and shadow
 - wind and sea surface condition
 - nature of seabed/sea bottom albedo
 - absorption and diffusion of light by the water column (turbidity, algae etc)
 - atmospheric condition
 - type of spectral bands









22–26 April, Hanoi, Vietnam





Background

Algorithms used for depth estimation/derivation from satellite images

- Banny and Dawson
 - Basic method.
 - Uses one band only.
- Lyzenga's
 - Uses multiple visible bands
 - Applicable to variable seafloor reflectivity
- Stumpf and Holderied
 - Adapting blue and green bands
- Lee's
 - Inversion method
 - Requires multiple bands (>5) and high resolution










22–26 April, Hanoi, Vietnam







SDB General Process Flow

- Geometric Correction
 - Correcting distortions of image
 - Improving horizontal accuracy
- Radiometric Correction
 - Calibration of recorded variance value reflected
 - Conversion to ToA radiance, ToA reflectance, sun-glint correction and atmospheric correction
- Radiometric Analysis for Depth Estimation
 - Ground truthing (radiative transfer model)
- Tidal Correction








THE SCIENCE OF WHERE





22-26 April, Hanoi, Vietnam





Study Motivation

Recognised as a cost and time effective solution - remote sensing environments Majority of the research conducted in Europe and American continents Few studies were carried out in Oceania and African coastal - outside of tropical climate environment Minimum sample of multispectral satellite images with limited study area Timely to have a detailed research in Malaysia - tropical environment No further analysis with survey standard - SP-44










22–26 April, Hanoi, Vietnam





The Study - Questions

How reliable this SDB technology in producing a consistent level of accuracy in the field of hydrographic surveying, especially for the shallow water area?

Which class of survey standard (IHO) that this technology capable of achieving?

What improvement can be developed to simplify the current processing procedure which can adopt into hydrographic surveying application?









22–26 April, Hanoi, Vietnam





The Study - Objectives

Examine & Identify the level of reliability produced from the algorithms focusing on the shallow water area which reflects on the hydrographic mapping applications

Analyse the degree of accuracy and standards of bathymetric data derived from SDB technology in meeting the standard and specifications laid by IHO

Improve the SDB empirical method results with optimisation technique focusing on the tropical environmental parameter











22–26 April, Hanoi, Vietnam





ORGANISED BY






PLATINUM SPONSORS








22-26 April, Hanoi, Vietnam






Research Area















22-26 April, Hanoi, Vietnam





Data

| Satellite Data/ Category | Spatial Resolution | Collection/ Owner | Area/Date Captured | Metadata |
|--------------------------------------|--------------------|---|--|--|
| Sentinel-2A (High-Resolution) | 10 m | Free download from ESA Website | Tawau (29 Nov 2016) Labuan (28 June 2017) | MSI_20161129T134807 MSI_20170628T094035 |
| Pleiades (Very High- Resolution) | 2 m | Supplied by Malaysian Remote Sensing Agency under cooperation with the National Hydrographic Centre | Tawau (12 July 2016) | DS_PHR1A_201607120241209 |
| WorldView2 (Very High-Resolution) | 0.5 m | Supplied by Universiti Malaysia Terengganu under joint research Marine Ecosystem Mapping for Taman Laut Labuan Project under Department of Marine Park Malaysia | Labuan (26 June 2015) | 15JUN26025812-P2AS- 056663834010 |










22-26 April, Hanoi, Vietnam





Data

| Bathymetry Data | Depth Sensor | Positioning System | Spatial Resolution/Survey Order | Survey Area |
|--------------------------------|--------------------|--------------------|------------------------------------|---------------|
| SBES (30 Nov to 2 Dec 2016) | Odom Mk III | Veripos LD4 | 0.5 m Special Order | Tawau Port |
| MBES (6 to 27 April 2017) | Kongsberg EM 2040C | C-Nav 3050 | 0.5 m Special Order | Pulau Kuraman |
| SBES (6 to 27 April 2017) | Odom Mk III | C-Nav 3050 | 0.5 m Special Order | Pulau Kuraman |











22-26 April, Hanoi, Vietnam





The Study-Processing (Tawau)












22-26 April, Hanoi, Vietnam





Processing (Pulau Kuraman)



Training Dataset

Full Dataset












22-26 April, Hanoi, Vietnam





Data Processing
















22-26 April, Hanoi, Vietnam





Depth Derivation









THE SCIENCE OF WHERE




22-26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"



Data Analysis












22-26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"



Model Enhancement












22-26 April, Hanoi, Vietnam





Model Enhancement












22-26 April, Hanoi, Vietnam





Model Enhancement












22–26 April, Hanoi, Vietnam





The Study- Model Enhancement





ORGANISED BY












22-26 April, Hanoi, Vietnam





Model Enhancement

 Apply simple filtering algorithm using least square adjustment method for linear regression and the coefficient of determination value formula

$$\widehat{X} = (A^T P A)^{-1} (A^T P F)$$

$$F = Ax$$

$$R^{2} = \frac{\sum (\widehat{y}_{i} - \overline{y})^{2}}{\sum (y_{i} - \overline{y})^{2}}$$












22-26 April, Hanoi, Vietnam















22-26 April, Hanoi, Vietnam



"Geospatial Information for a Smarter Life and Environmental Resilience"

| Regression Summary (Stumpf Tawau Port) | | | | | |
|--|----------|----------|--|--|--|
| Sentinel-2A Pleiades | | | | | |
| R Square | 0.5948 | 0.6214 | | | |
| Slope (Gain) | 321.32 | 427.52 | | | |
| Y-Axis Intercept (Offset) | - 326.85 | - 383.49 | | | |
| Observations | 196 | 199 | | | |

| Regressions Summary (M-LOT Tawau Port) | | | | | | |
|--|---------------|-------------|-------------|---------------|-------------|-------------|
| | | Sentinel-2A | | Pleiades | | |
| | Single Linear | Lower Layer | Upper Layer | Single Linear | Lower Layer | Upper Layer |
| R Square | 0.7276 | 0.6035 | 0.1943 | 0.7424 | 0.3663 | 0.4761 |
| Slope (Gain) | 321.55 | 314.31 | 282.0 | 460.33 | 385.28 | 373.08 |
| Y-Axis Intercept (Offset) | -327.11 | -319.72 | -285.64 | -413.61 | -345.93 | -333.08 |
| Observations | 166 | 84 | 86 | 183 | 88 | 95 |












22-26 April, Hanoi, Vietnam


























22-26 April, Hanoi, Vietnam







| Regression Summary (Stumpf Pulau Kuraman) | | | | | |
|---|---------|---------|--|--|--|
| Sentinel-2A WorldView-2 | | | | | |
| R Square | 0.5665 | 0.4326 | | | |
| Slope (Gain) | 211.4 | 217.29 | | | |
| Y-Axis Intercept (Offset) | -212.69 | -179.03 | | | |
| Observations | 492 | 496 | | | |

| Regressions Summary (M-LOT Pulau Kuraman) | | | | | | |
|---|---------------|-------------|-------------|---------------|-------------|-------------|
| | Sentinel-2A | | | WorldView-2 | | |
| | Single Linear | Lower Layer | Upper Layer | Single Linear | Lower Layer | Upper Layer |
| R Square | 0.5968 | 0.6798 | 0.5626 | 0.4326 | 0.7544 | 0.6055 |
| Slope (Gain) | 206.17 | 68.716 | 305.65 | 217.29 | 84.056 | 1295.7 |
| Y-Axis Intercept (Offset) | -207.34 | -67.634 | 310 | -179.03 | -67.894 | -1095.3 |
| Observations | 488 | 165 | 328 | 492 | 225 | 272 |













22-26 April, Hanoi, Vietnam






Results & Analysis







Lyzenga model

Stumpf model

M-LOT model













22-26 April, Hanoi, Vietnam






Results & Analysis







Lyzenga model

Stumpf model

M-LOT model














22-26 April, Hanoi, Vietnam











Lyzenga model

Stumpf model

M-LOT model
















22-26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"









Lyzenga model

Stumpf model

M-LOT model












22-26 April, Hanoi, Vietnam





| | Sentinel-2A (Tawau Port) | | Pleaides (Tawau Port) | | t) | |
|--------------------|--------------------------|-----------|-----------------------|-----------|----------|----------|
| | Lyzenga | Stumpf | M-LOT | Lyzenga | Stumpf | M-LOT |
| Sample Variance | 5.159 | 5.124 | 4.713 | 3.992 | 3.963 | 3.240 |
| RMSE | 4.464 | 4.459 | 4.307 | 4.323 | 4.435 | 4.379 |
| Standard Deviation | 2.271 | 2.264 | 2.171 | 1.998 | 1.991 | 1.800 |
| Sum | -2203.843 | -2196.262 | -1409.845 | -1061.813 | -917.150 | -272.315 |
| Sample Count | 2113 | 2113 | 2113 | 2213 | 2213 | 2213 |










22-26 April, Hanoi, Vietnam





| | Sentinel-2A (Pulau Kuraman) | | WV | 2 (Pulau Kuran | nan) | |
|--------------------|-----------------------------|----------|------------|----------------|-------------|------------|
| | Lyzenga | Stumpf | M-LOT | R-Lyzenga | Stumpf | M-LOT |
| Sample Variance | 4.393 | 5.813 | 5.105 | 2.676 | 3.887 | 3.437 |
| RMSE | 2.118 | 2.411 | 2.263 | 1.841 | 2.975 | 1.958 |
| Standard Deviation | 2.096 | 2.411 | 2.259 | 1.634 | 1.972 | 1.854 |
| Sum | -79793.673 | 9136.311 | -32656.619 | 424317.754 | 1118489.646 | 316645.097 |
| Sample Count | 259013 | 259013 | 259013 | 502081 | 502081 | 502081 |










22-26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"



| | Sentinel-2A (Tawau Port) | | | | |
|----------------------|--------------------------|--------------|-----------|--|--|
| | Lyzenga | Stumpf | M-LOT | | |
| Total Samples | 2113 | 2113 | 2113 | | |
| IHO Passed | 822 | 826 | 906 | | |
| IHO Failed | 1291 | 1287 | 1207 | | |
| | IHO Surv | ey Order Dis | tribution | | |
| Special Order | 209 | 207 | 198 | | |
| Order 1A/1B | 297 | 299 | 399 | | |
| Order 2 | 316 | 320 | 309 | | |












22-26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"



| | Sentinel-2A (Tawau Port) | | | | |
|---------------|--------------------------|--------------|-----------|--|--|
| | Lyzenga | Stumpf | M-LOT | | |
| Total Samples | 2113 | 2113 | 2113 | | |
| IHO Passed | 822 | 826 | 906 | | |
| IHO Failed | 1291 | 1287 | 1207 | | |
| | IHO Surv | ey Order Dis | tribution | | |
| Special Order | 209 | 207 | 198 | | |
| Order 1A/1B | 297 | 299 | 399 | | |
| Order 2 | 316 | 320 | 309 | | |















22-26 April, Hanoi, Vietnam





| | Pleiades (Tawau Port) | | | | |
|----------------------|-------------------------------|--------|-------|--|--|
| | Lyzenga | Stumpf | M-LOT | | |
| Total Samples | 2213 | 2213 | 2213 | | |
| IHO Passed | 1019 | 1016 | 1149 | | |
| IHO Failed | 1194 | 1197 | 1064 | | |
| | IHO Survey Order Distribution | | | | |
| Special Order | 214 | 217 | 283 | | |
| Order 1A/1B | 417 | 429 | 500 | | |
| Order 2 | 388 | 370 | 366 | | |

















22-26 April, Hanoi, Vietnam





| | Pleiades (Tawau Port) | | | | |
|---------------|-----------------------|--------------|------------|--|--|
| | Lyzenga | Stumpf | M-LOT | | |
| Total Samples | 2213 | 2213 | 2213 | | |
| IHO Passed | 1019 | 1016 | 1149 | | |
| IHO Failed | 1194 | 1197 | 1064 | | |
| | IHO Surve | ey Order Dis | stribution | | |
| Special Order | 214 | 217 | 283 | | |
| Order 1A/1B | 417 | 429 | 500 | | |
| Order 2 | 388 | 370 | 366 | | |












22-26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"



| | Sentinel2A (Pulau Kuraman) | | |
|---------------|-------------------------------|--------|--------|
| | Lyzenga | Stumpf | M-LOT |
| Total Samples | 460252 | 460252 | 460252 |
| IHO Passed | 166747 | 103418 | 151100 |
| IHO Failed | 293505 | 356834 | 309152 |
| | IHO Survey Order Distribution | | |
| Special Order | 34245 | 18401 | 32311 |
| Order 1A/1B | 67143 | 39617 | 61991 |
| Order 2 | 65359 | 45400 | 56798 |















22-26 April, Hanoi, Vietnam





| | Sentinel2A (Pulau Kuraman) | | |
|---------------|-------------------------------|--------|--------|
| | Lyzenga | Stumpf | M-LOT |
| Total Samples | 460252 | 460252 | 460252 |
| IHO Passed | 166747 | 103418 | 151100 |
| IHO Failed | 293505 | 356834 | 309152 |
| | IHO Survey Order Distribution | | |
| Special Order | 34245 | 18401 | 32311 |
| Order 1A/1B | 67143 | 39617 | 61991 |
| Order 2 | 65359 | 45400 | 56798 |















22-26 April, Hanoi, Vietnam





| | Worldview-2 (Pulau Kuraman) | | |
|---------------|-------------------------------|--------|--------|
| | Lyzenga | Stumpf | M-LOT |
| Total | 502081 | 502081 | 502081 |
| Samples | | | |
| IHO Passed | 239639 | 123147 | 235260 |
| IHO Failed | 262442 | 378934 | 266821 |
| | IHO Survey Order Distribution | | |
| Special Order | 50692 | 14364 | 50167 |
| Order 1A/1B | 98634 | 43408 | 98235 |
| Order 2 | 90313 | 65375 | 86858 |














22-26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"



| | Worldview-2 (Pulau Kuraman) | | |
|---------------|-------------------------------|--------|--------|
| | Lyzenga | Stumpf | M-LOT |
| Total Samples | 502081 | 502081 | 502081 |
| IHO Passed | 239639 | 123147 | 235260 |
| IHO Failed | 262442 | 378934 | 266821 |
| | IHO Survey Order Distribution | | |
| Special Order | 50692 | 14364 | 50167 |
| Order 1A/1B | 98634 | 43408 | 98235 |
| Order 2 | 90313 | 65375 | 86858 |











22-26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"



Difficult to have good quality image

Minimum change to have an ideal condition/'acceptable' quality

Complex processing

- SDB for hydrography application heavily required an experience hydrographic surveyor supervision
- Tedious calibration techniques

Rough replication of seabed

Features not always detected and/or difficult to determine

Low depth (acceptable) penetration

- 10 m on average depends on water clarity
- Unable to meet S-44 Special Order and Order 1a

Sea bottom albedo effect

- Radiance/luminance very depending on seabed types, alga etc
- Require detail analysis especially on the factor of bottom type/albedo









22-26 April, Hanoi, Vietnam





SDB is not a rhetorical application for shallow water area in tropical environment

• Further study still required to improve the consistency (very shallow area)

Urgent need - to establish the standards for SDB application

Achievable as the new IHO SP-44 will have entirely new concept/objective oriented

About time to have a detail guideline for SDB application (shallow water area)

Adopting this promising technology as an assessment tool

Positive progress and promising results

Will give significant contributions to HOs/relevant agencies

More area to analyse especially on the factor of bottom type/albedo









22-26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"



Dedication


Special thanks to various organisations and agencies for the

unconditional support to this study especially to

National Hydrographic Centre

Department of Survey and Mapping Malaysia

- Malaysia Remote Sensing Agency
- Board of Land Surveyors Malaysia
- Audience thank you for chook











22-26 April, Hanoi, Vietnam







chúc một ngày tốt lành





